

SK-TTC → Genes

DNA sequence 1600 b.p. ggaacagctat ... gtcgttttata linear

1 ggaacagctatgaccatgattacgccaagctcgaaattaccctcactaaagggaacaaagctggagctcggtaccg 80

81 ggccacc ATG GTT TTT TCA ACA CCA ATT CCA TTT TCT TAT TCT AAA AAT CTG GAT TGT TGG 141
 1 M V F S T P I P F S Y S K N L D C W 18

142 GTT GAT AAT GAA GAA GAT ATA GAT GTT ATA TTA AAA AAG AGT ACA ATT TTA AAT TTA GAT 201
 19 V D N E E D I D V I L K K S T I L N L D 38

202 ATT AAT AAT GAT ATT ATA TCA GAT ATA TCT GGG TTT AAT TCA TCT GTA ATA ACA TAT CCA 261
 39 I N N D I I S D I S G F N S S V I T Y P 58

262 GAT GCT CAA TTG GTG CCC GGA ATA AAT GGC AAA GCA ATA CAT TTA GTA AAC AAT GAA TCT 321
 59 D A Q L V P G I N G A I H L V N N E S 78

322 TCT GAA GTT ATA GTG CAT AAA GCT ATG GAT ATT GAA TAT AAT GAT ATG TTT AAT AAT TTT 381
 79 S E V I V H K A M D I E Y N D M F N N F 98

382 ACC GTT AGC TTT TGG TTG AGG GTT CCT AAA GTA TCT GCT AGT CAT TTA GAA CAA TAT GGC 441
 99 T V S F W L R V P K V S A S H L E Q Y G 118

442 ACA AAT GAG TAT TCA ATA ATT AGC TCT ATG AAA AAA CAT AGT CTA TCA ATA GGA TCT GGT 501
 119 T N E Y S I I S S M K K H S L S I G S G 138

502 TGG AGT GTA TCA CTT AAA GGT AAT AAC TTA ATA TGG ACT TTA AAA GAT TCC GCG GGA GAA 561
 139 W S V S L K G N N L I I W T L K D S A G E 158

562 GTT AGA CAA ATA ACT TTT AGG GAT TTA CCT GAT AAA TTT AAT GCT TAT TTA GCA AAT AAA 621
 159 V R Q I T F R D L P D K F N A Y L A N K 178

622 TGG GTT TTT ATA ACT ATT ACT AAT GAT AGA TTA TCT TCT GCT AAT TTG TAT ATA AAT GGA 681
 179 W V F I T I T N D R L S S A N L Y I N G 198

682 GTA CTT ATG GGA AGT GCA GAA ATT ACT GGT TTA GGA GCT ATT AGA GAG GAT AAT AAT ATA 741
 199 V L M G S A E I T G L G A I R E D N N I 218

FIG. 1A

742 ACA TTA AAA CTA GAT AGA TGT AAT AAT AAT AAT CAA TAC GTT TCT ATT GAT AAA TTT AGG 801
219 T L K L D R C N N N N Q Y V S I D K F R 238

802 ATA TTT TGC AAA GCA TTA AAT CCA AAA GAG ATT GAA AAA TTA TAC ACA AGT TAT TTA TCT 861
239 I F C K A L N P K E I E K L Y T S Y L S 258

862 ATA ACC TTT TTA AGA GAC TTC TGG GGA AAC CCT TTA CGA TAT GAT ACA GAA TAT TAT TTA 921
259 I T F L R D F W G N P L R Y D T E Y Y L 278

922 ATA CCA GTA GCT TCT AGT TCT AAA GAT GTT CAA TTG AAA AAT ATA ACA GAT TAT ATG TAT 981
279 I P V A S S S K D V Q L K N I T D Y M Y 298

982 TTG ACA AAT GCG CCA TCG TAT ACT AAC GGA AAA TTG AAT ATA TAT TAT AGA AGG TTA TAT 1041
299 L T N A P S Y T N G K L N I Y Y R R L Y 318

1042 AAT GGA CTA AAA TTT ATT ATA AAA AGA TAT ACA CCT AAT AAT AAT GAA ATA GAT TCT TTT GTT 1101
319 N G L K F I I K R Y T P N N E I D S F V 338

1102 AAA TCA GGT GAT TTT ATT AAA TTA TAT GTA TCA TAT AAC AAT AAT GAG CAC ATT GTA GGT 1161
339 K S G D F I K L Y V S Y N N N E H I V G 358

1162 TAT CCG AAA GAT GGA AAT GCC TTT AAT AAT AAT CTT GAT AGA ATT CTA AGA GTA GGT TAT AAT 1221
359 Y P K D G N A F N N L D R I L R V G Y N 378

1222 GCC CCA GGT ATC CCT CTT TAT AAA AAT ATG GAA GCA GTA AAA TTG CGT GAT TTA AAA ACC 1281
379 A P G I P L Y K K M E A V K L R D L K T 398

1282 TAT TCT GTA CAA CTT AAA TTA TAT GAT GAT AAA AAT GCA TCT TTA GGA CTA GTA GGT ACC 1341
399 Y S V Q L K L Y D D K N A S L G L V G T 418


1342 CAT AAT GGT CAA ATA GGC AAC GAT CCA AAT AGG GAT ATA TTA ATT GCA AGC AAC TGG TAC 1401
419 H N G Q I G N D P N R D I L I A S N W Y 438

1402 TTT AAT CAT TTA AAA GAT AAA ATT TTA GGA TGT GAT TGG TAC TTT GTA CCT ACA GAT GAG 1461
439 F N H L K D K I L G C D W Y F V P T D E 458

1462 GGA TGG ACA AAT GAT TAA acagattgatattgttcacatattgccccgggattcctctagagtcgacctcgaggg 1535
459 G W T N D * (SEQ ID NO:2) 464

1536 ggggccccgtaccaccaattgcgcctatagtgagtcgtattacaattcactggccgctgctttttacaa (SEQ ID NO:1) 1600

FIG. 1B



TTC cDNA isolation:

The TTC cDNA was isolated from a *Clostridium Tetani* strain using Polymerase Chain Reaction. We used a three times PCR to generate three overlapping fragments respectively of 465 bp (PCR1; primer 1: 5'-CCC CCC GGG CCA CCA TGG TTT TTT CAA CAC CAA TTC CAT TTT CTT ATT C-3' (SEQ ID NO:4) & primer 2: 5'-CTA AAC CAG TAA TTT CTG-3' (SEQ ID NO:5), of 648 bp (PCR2; primer 3: 5'-AAT TAT GGA CTT TAA AAG ATT CCG C-3' (SEQ ID NO:6) & primer 4: 5'-GGC ATT ATA ACC TAC TCT TAG AAT-3' (SEQ ID NO:7) and of 338 bp (PCR3; primer 5: 5'-AAT GCC TTT AAT AAT CTT GAT AGA AAT-3' (SEQ ID NO:8) & primer 6: 5'-CCC CCC GGG CAT ATG TCA TGA ACA TAT CAA TCT GTT TAA TC-3' (SEQ ID NO:9), and each fragment was sequentially cloned into pBluescript KS+ to produce plasmid pBS-TTC. The upstream primer 1 contained the Ribosome Binding Site (RBS) and translation initiation signals.

FIG. 2




FIG. 3

09816467-062901




FIG. 4

09816467 062901

pCMV.LACZ.TTC -> Graphic Map
 DNA sequence 8519 b.p. tagttattaata ... accgccatgcat circular




FIG. 5

09816467.062901